

TOPICAL PAST PAPER QUESTIONS WORKBOOK

AS & A Level Mathematics (9709) Paper 1 [Pure Mathematics 1]

May/June 2015 – February/March 2022

Chapter 8

Integration

- (c) Find $\frac{d^2y}{dx^2}$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (d) Determine the nature of the stationary point at $(2, -3.5)$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Find the coordinates of the stationary points on the curve. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Find $f''(x)$. [1]

.....

.....

.....

.....

- (d) Hence, or otherwise, determine the nature of each of the stationary points. [2]

.....

.....

.....

.....

.....

.....

451. 9709_w21_qp_11 Q: 10

(a) Find $\int_1^{\infty} \frac{1}{(3x-2)^{\frac{3}{2}}} dx$. [4]

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



The diagram shows the curve with equation $y = \frac{1}{(3x-2)^{\frac{3}{2}}}$. The shaded region is bounded by the curve, the x -axis and the lines $x = 1$ and $x = 2$. The shaded region is rotated through 360° about the x -axis.

(b) Find the volume of revolution. [4]

.....

.....

.....


.....

.....

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

Dotted lines for writing.

463. 9709_w20_qp_11 Q: 12



The diagram shows a curve with equation $y = 4x^{\frac{1}{2}} - 2x$ for $x \geq 0$, and a straight line with equation $y = 3 - x$. The curve crosses the x-axis at $A(4, 0)$ and crosses the straight line at B and C .

- (a) Find, by calculation, the x-coordinates of B and C . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Show that B is a stationary point on the curve. [2]

.....

.....

.....

.....

.....

.....

.....

472. 9709_s19_qp_12 Q: 3

A curve is such that $\frac{dy}{dx} = x^3 - \frac{4}{x^2}$. The point $P(2, 9)$ lies on the curve.

- (i) A point moves on the curve in such a way that the x -coordinate is decreasing at a constant rate of 0.05 units per second. Find the rate of change of the y -coordinate when the point is at P . [2]

.....
.....
.....
.....
.....
.....
.....
.....

- (ii) Find the equation of the curve. [3]

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....



- (ii) Find the coordinates of M . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The shaded region is bounded by the curve, the y -axis and the line through M parallel to the x -axis.

- (iii) Find, showing all necessary working, the area of the shaded region. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



(ii) Find the coordinates of B .

[4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Find, showing all necessary working, the area of the shaded region.

[4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



- (ii) A point P moves along the curve in such a way that the y -coordinate is increasing at a constant rate of 0.06 units per second. Find the rate of change of the x -coordinate when P passes through $(2, 5)$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (iii) Show that $\frac{d^2y}{dx^2} \times \frac{dy}{dx}$ is constant. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Determine, showing all necessary working, the nature of the stationary point. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



(ii) Find the coordinates of M . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Find, showing all necessary working, the area of the shaded region. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....



506. 9709_m16_qp_12 Q: 2

A curve for which $\frac{dy}{dx} = 3x^2 - \frac{2}{x^3}$ passes through $(-1, 3)$. Find the equation of the curve. [4]

507. 9709_m16_qp_12 Q: 10




The diagram shows part of the curve $y = \frac{1}{16}(3x - 1)^2$, which touches the x -axis at the point P . The point $Q(3, 4)$ lies on the curve and the tangent to the curve at Q crosses the x -axis at R .

- (i) State the x -coordinate of P . [1]

Showing all necessary working, find by calculation

- (ii) the x -coordinate of R , [5]
(iii) the area of the shaded region PQR . [6]
-

508. 9709_s16_qp_11 Q: 3



The diagram shows part of the curve $x = \frac{12}{y^2} - 2$. The shaded region is bounded by the curve, the y-axis and the lines $y = 1$ and $y = 2$. Showing all necessary working, find the volume, in terms of π , when this shaded region is rotated through 360° about the y-axis. [5]

509. 9709_s16_qp_11 Q: 4

A curve is such that $\frac{dy}{dx} = 2 - 8(3x + 4)^{-\frac{1}{2}}$.

- (i) A point P moves along the curve in such a way that the x -coordinate is increasing at a constant rate of 0.3 units per second. Find the rate of change of the y -coordinate as P crosses the y -axis. [2]


The curve intersects the y -axis where $y = \frac{4}{3}$.

- (ii) Find the equation of the curve. [4]
-

510. 9709_s16_qp_12 Q: 2

A curve is such that $\frac{dy}{dx} = \frac{8}{(5-2x)^2}$. Given that the curve passes through (2, 7), find the equation of the curve. [4]


511. 9709_s16_qp_12 Q: 10



The diagram shows the part of the curve $y = \frac{8}{x} + 2x$ for $x > 0$, and the minimum point M .

- (i) Find expressions for $\frac{dy}{dx}$, $\frac{d^2y}{dx^2}$ and $\int y^2 dx$. [5]
- (ii) Find the coordinates of M and determine the coordinates and nature of the stationary point on the part of the curve for which $x < 0$. [5]
- (iii) Find the volume obtained when the region bounded by the curve, the x -axis and the lines $x = 1$ and $x = 2$ is rotated through 360° about the x -axis. [2]
-

512. 9709_s16_qp_13 Q: 2



The diagram shows part of the curve $y = (x^3 + 1)^{\frac{1}{2}}$ and the point $P(2, 3)$ lying on the curve. Find, showing all necessary working, the volume obtained when the shaded region is rotated through 360° about the x -axis. [4]


513. 9709_s16_qp_13 Q: 3

A curve is such that $\frac{dy}{dx} = 6x^2 + \frac{k}{x^3}$ and passes through the point $P(1, 9)$. The gradient of the curve at P is 2.

(i) Find the value of the constant k . [1]

(ii) Find the equation of the curve. [4]

514. 9709_w16_qp_11 Q: 7



The diagram shows parts of the curves $y = (2x - 1)^2$ and $y^2 = 1 - 2x$, intersecting at points A and B .

- (i) State the coordinates of A . [1]
- (ii) Find, showing all necessary working, the area of the shaded region. [6]
-

515. 9709_w16_qp_11 Q: 10

A curve has equation $y = f(x)$ and it is given that $f'(x) = 3x^{\frac{1}{2}} - 2x^{-\frac{1}{2}}$. The point A is the only point on the curve at which the gradient is -1 .

- (i) Find the x -coordinate of A . [3]
- (ii) Given that the curve also passes through the point $(4, 10)$, find the y -coordinate of A , giving your answer as a fraction. [6]
-

516. 9709_w16_qp_12 Q: 1

A curve is such that $\frac{dy}{dx} = \frac{8}{\sqrt{4x+1}}$. The point (2, 5) lies on the curve. Find the equation of the curve. [4]

517. 9709_w16_qp_13 Q: 10

A curve is such that $\frac{dy}{dx} = \frac{2}{a}x^{-\frac{1}{2}} + ax^{-\frac{3}{2}}$, where a is a positive constant. The point $A(a^2, 3)$ lies on the curve. Find, in terms of a ,

- (i) the equation of the tangent to the curve at A , simplifying your answer, [3]
(ii) the equation of the curve. [4]

It is now given that $B(16, 8)$ also lies on the curve.


- (iii) Find the value of a and, using this value, find the distance AB . [5]
-

518. 9709_w16_qp_13 Q: 11

A curve has equation $y = (kx - 3)^{-1} + (kx - 3)$, where k is a non-zero constant.


- (i) Find the x -coordinates of the stationary points in terms of k , and determine the nature of each stationary point, justifying your answers. [7]

(ii)



The diagram shows part of the curve for the case when $k = 1$. Showing all necessary working, find the volume obtained when the region between the curve, the x -axis, the y -axis and the line $x = 2$, shown shaded in the diagram, is rotated through 360° about the x -axis. [5]

519. 9709_s15_qp_11 Q: 10



The diagram shows part of the curve $y = \frac{8}{\sqrt{3x+4}}$. The curve intersects the y -axis at $A(0, 4)$. The normal to the curve at A intersects the line $x = 4$ at the point B .

(i) Find the coordinates of B . [5]

(ii) Show, with all necessary working, that the areas of the regions marked P and Q are equal. [6]

520. 9709_s15_qp_12 Q: 10


The equation of a curve is $y = \frac{4}{2x-1}$.

- (i) Find, showing all necessary working, the volume obtained when the region bounded by the curve, the x -axis and the lines $x = 1$ and $x = 2$ is rotated through 360° about the x -axis. [4]
- (ii) Given that the line $2y = x + c$ is a normal to the curve, find the possible values of the constant c . [6]

521. 9709_s15_qp_13 Q: 2

A curve is such that $\frac{dy}{dx} = (2x + 1)^{\frac{1}{2}}$ and the point $(4, 7)$ lies on the curve. Find the equation of the curve. [4]

522. 9709_s15_qp_13 Q: 10



Points $A(2, 9)$ and $B(3, 0)$ lie on the curve $y = 9 + 6x - 3x^2$, as shown in the diagram. The tangent at A intersects the x -axis at C . Showing all necessary working,


- (i) find the equation of the tangent AC and hence find the x -coordinate of C , [4]
- (ii) find the area of the shaded region ABC . [5]
-

523. 9709_w15_qp_11 Q: 2

The function f is such that $f'(x) = 3x^2 - 7$ and $f(3) = 5$. Find $f(x)$.

[3]

524. 9709_w15_qp_11 Q: 11




The diagram shows part of the curve $y = (1 + 4x)^{\frac{1}{2}}$ and a point $P(6, 5)$ lying on the curve. The line PQ intersects the x -axis at $Q(8, 0)$.

- (i) Show that PQ is a normal to the curve. [5]
- (ii) Find, showing all necessary working, the exact volume of revolution obtained when the shaded region is rotated through 360° about the x -axis. [7]

[In part (ii) you may find it useful to apply the fact that the volume, V , of a cone of base radius r and vertical height h , is given by $V = \frac{1}{3}\pi r^2 h$.]

525. 9709_w15_qp_12 Q: 10



The diagram shows part of the curve $y = \sqrt{9 - 2x^2}$. The point $P(2, 1)$ lies on the curve and the normal to the curve at P intersects the x -axis at A and the y -axis at B .

- (i) Show that B is the mid-point of AP . [6]

The shaded region is bounded by the curve, the y -axis and the line $y = 1$.

- (ii) Find, showing all necessary working, the exact volume obtained when the shaded region is rotated through 360° about the y -axis. [5]

526. 9709_w15_qp_13 Q: 9


A curve passes through the point $A(4, 6)$ and is such that $\frac{dy}{dx} = 1 + 2x^{-\frac{1}{2}}$. A point P is moving along the curve in such a way that the x -coordinate of P is increasing at a constant rate of 3 units per minute.

- (i) Find the rate at which the y -coordinate of P is increasing when P is at A . [3]
 - (ii) Find the equation of the curve. [3]
 - (iii) The tangent to the curve at A crosses the x -axis at B and the normal to the curve at A crosses the x -axis at C . Find the area of triangle ABC . [5]
-

527. 9709_w15_qp_13 Q: 10

The function f is defined by $f(x) = 2x + (x + 1)^{-2}$ for $x > -1$.

- (i) Find $f'(x)$ and $f''(x)$ and hence verify that the function f has a minimum value at $x = 0$. [4]



The points $A(-\frac{1}{2}, 3)$ and $B(1, 2\frac{1}{4})$ lie on the curve $y = 2x + (x + 1)^{-2}$, as shown in the diagram.

- (ii) Find the distance AB . [2]
- (iii) Find, showing all necessary working, the area of the shaded region. [6]
-

